arXiv:1207.7165v1 [math.NT] 31 Jul 2012

SOME IDENTITIES INVOLVING GEGENBAUER POLYNOMIALS

DAE SAN KIM[1], TAEKYUN KIM[2], AND SEOG-HOON RIM[3]

Abstract. In this paper we derive some interesting identities involving Gegenbauer polynomials arising from the orthogonality of Gegenbauer polynomials for the inner product space $\mathbb{P}_n$ with respect to the weighted inner product $< p_1, p_2 >= \int_{-1}^{1} p_1(x)p_2(x)(1-x^2)^{\lambda-\frac{1}{2}}dx$.

1. Introduction

The Gegenbauer polynomials are given in terms of the Jacobi polynomials $P_n^{(\alpha,\beta)}(x)$ with $\alpha=\beta=\lambda-\frac{1}{2}$ $(\lambda>-\frac{1}{2},\ \lambda\neq 0)$ by

$$\begin{aligned} C_n^{(\lambda)}(x) &= \frac{\Gamma\left(\lambda+\frac{1}{2}\right)\Gamma\left(n+2\lambda\right)}{\Gamma(2\lambda)\Gamma\left(n+\lambda+\frac{1}{2}\right)}P_n^{\left(\lambda-\frac{1}{2},\lambda-\frac{1}{2}\right)}(x) \\ &= \binom{n+2\lambda-1}{n}\sum_{k=0}^{n}\frac{\binom{n}{k}(2\lambda+n)_k}{\left(\lambda+\frac{1}{2}\right)_k}\left(\frac{x-1}{2}\right)^k \end{aligned} \tag{1.1}$$

where $(a)_k = a(a+1)(a+2)\cdots(a+k-1)$, (see [19, 23]).

From (1.1), we note that $C_k^{(\lambda)}(x)$ is a polynomial of degree n with real coefficients and $C_n^{(\lambda)}(1)=\binom{n+2\lambda-1}{n}$. The leading coefficient of $C_n^{(\lambda)}(x)$ is $2^n\binom{\lambda+n-1}{n}$. By the theory of Jacobi polynomials with $\alpha=\beta=\lambda-\frac{1}{2}$, $\lambda>-\frac{1}{2}$, and $\lambda\neq 0$, we get

$$C_n^{(\lambda)}(-x) = (-1)^n C_n^{(\lambda)}(x). \tag{1.2}$$

It is not difficult to show that $C_n^{(\lambda)}(x)$ is a solution of the following Gegenbauer differential equation:

$$(1-x^2)y^{''} - (2\lambda+1)xy^{'} + n(n+2\lambda)y = 0.$$

The Rodrigues' formula for the Gegenbauer polynomials are well known as the following:

$$(1-x^2)^{\lambda-\frac{1}{2}}C_n^{(\lambda)}(x) = \frac{(-2)^n(\lambda)_n}{n!(n+2\lambda)_n}\left(\frac{d}{dx}\right)^n(1-x^2)^{n+\lambda-\frac{1}{2}},\ \text{(see [1, 13])}. \tag{1.3}$$

The equation (1.3) can be easily derived from the properties of Jacobi polynomials.

As is well known, the generating function of Gegenbauer polynomials is given by

$$\frac{2^{\lambda-\frac{1}{2}}}{(1-2xt+t^2)^{\frac{1}{2}}(1-xt+\sqrt{1-2xt+t^2})^{\lambda-\frac{1}{2}}} = \sum_{n=0}^{\infty}\frac{\left(\lambda+\frac{1}{2}\right)_n}{(2\lambda)_n}C_n^{(\lambda)}(x)t^n. \tag{1.4}$$

The equation (1.4) can be also derived from the generating function of Jacobi polynomials.

From (1.4), we note that

$$\frac{1}{(1-2xt+t^2)^\lambda}=\sum_{n=0}^{\infty}C_n^{(\lambda)}(x)t^n,\ (|t|<1,\ |x|\leq 1). \tag{1.5}$$

The proof of (1.5) is given in the following book: Stein & Weiss, Introduction to Fourier Analysis in Euclidean space, Princeton University Press, 1971.

By (1.1) and (1.2), we get

$$\int_{-1}^{1}C_n^{(\lambda)}(x)C_m^{(\lambda)}(x)(1-x^2)^{\lambda-\frac{1}{2}}dx=\frac{\pi 2^{1-2\lambda}\Gamma(n+2\lambda)}{n!(n+\lambda)(\Gamma(\lambda))^2}\delta_{m,n} \tag{1.6}$$

where $\delta_{m,n}$ is the Kronecker symbol and it holds for each fixed $\lambda\in\mathbb{R}$ with $\lambda>-\frac{1}{2}$ and $\lambda\neq 0$.

The equation (1.6) implies the orthogonality of $C_n^{(\lambda)}(x)$ and the equation (1.6) is important in deriving our results in this paper. From (1.5), we can derive the following derivative of Gehenbauer polynomials $C_n^{(\lambda)}(x)$:

$$\frac{d}{dx}C_n^{(\lambda)}(x)=2\lambda C_{n-1}^{(\lambda+1)}(x),\ \text{for } n\geq 1. \tag{1.7}$$

By (1.8), we get

$$\frac{d^k}{dx^k}C_n^{(\lambda)}(x)=2^k\lambda^kC_{n-k}^{(\lambda+k)}(x). \tag{1.8}$$

As is well known, the Bernoulli polynomials $B_n(x)$ are defined by the generating function to be

$$\frac{t}{e^t-1}e^{xt}=e^{B(x)t}=\sum_{n=0}^{\infty}B_n(x)\frac{t^n}{n!},\ \text{(see [2-10])}, \tag{1.9}$$

with the usual convention about replacing $B^n(x)$ by $B_n(x)$. In the special case, $x=0$, $B_n(0)=B_n$ are called the n-th Bernoulli numbers.

From (1.9), we note that

$$B_n(x)=(B+x)^n=\sum_{l=0}^{n}\binom{n}{l}B_{n-l}x^l,\ \text{(see [6-10])}, \tag{1.10}$$

and

$$B_n^{'}(x)=\frac{d}{dx}B_n(x)=nB_{n-1}(x). \tag{1.11}$$

The Euler polynomials $E_n(x)$ are also defined by the generating function to be

$$\frac{2}{e^t+1}e^{xt}=e^{E(x)t}=\sum_{n=0}^{\infty}E_n(x)\frac{t^n}{n!},\ \text{(see [12-17])}, \tag{1.12}$$

with the usual convention about replacing $E^n(x)$ by $E_n(x)$. In the special case, $x=0$, $E_n(0)=E_n$ are called the n-th Euler numbers. By (1.12), we see that the recurrence formula for E_n is given by

$$E_0=1,\ (E+1)^n+E_n=2\delta_{0,n},\ \text{(see [20-22])}. \tag{1.13}$$

For each fixed $\lambda\in\mathbb{R}$ with $\lambda>-\frac{1}{2}$ and $\lambda\neq 0$, let $\mathbb{P}_n=\{p(x)\in\mathbb{R}[x]\mid \deg p(x)\leq n\}$ be inner product space with respect to the inner product

$$<p_1(x),p_2(x)>=\int_{-1}^{1}(1-x^2)^{\lambda-\frac{1}{2}}p_1(x)p_2(x)dx, \tag{1.14}$$

where $p_1(x), p_2(x) \in \mathbb{P}_n$.

In this paper, we derive some interesting identities involving Gegenbauer polynomials arising from the orthogonality of those for the inner product space $\mathbb{P}_n$ with respect to the weighted inner product $< p_1, p_2 >= \int_{-1}^{1} p_1(x)p_2(x)(1-x^2)^{\lambda-\frac{1}{2}}dx$.

Our methods used in this paper are useful in finding some new identities and relations on the Bernoulli and Euler polynomials involving Gegenbauer polynomials.

2. Some identities involving Gegenabauer polynomials

Let us take $p(x) = \sum_{k=0}^{n} d_k C_k^{(\lambda)}(x) \in \mathbb{P}_n,\ d_k \in \mathbb{R}$. Then, by (1.6) and (1.14), we get

$$\begin{aligned} &< p(x), C_k^{(\lambda)}(x) >= d_k < C_k^{(\lambda)}(x), C_k^{(\lambda)}(x) > \\ &= d_k \int_{-1}^{1} (1-x^2)^{\lambda-\frac{1}{2}} C_k^{(\lambda)}(x) C_k^{(\lambda)}(x) dx = d_k \frac{\pi 2^{1-2\lambda}\Gamma(k+2\lambda)}{k!(k+\lambda)(\Gamma(\lambda))^2}. \end{aligned} \tag{2.1}$$

Thus, from (2.1), we have

$$d_k = \frac{(\Gamma(\lambda))^2 k!(k+\lambda)}{\pi 2^{1-2\lambda}\Gamma(k+2\lambda)} \int_{-1}^{1} (1-x^2)^{\lambda-\frac{1}{2}} p(x) C_k^{(\lambda)}(x) dx. \tag{2.2}$$

By (1.3) and (2.2), we get

$$\begin{aligned} d_k &= \frac{(\Gamma(\lambda))^2 k!(k+\lambda)}{\pi 2^{1-2\lambda}\Gamma(k+2\lambda)} \times \frac{(-2)^k(\lambda)_k}{k!(k+2\lambda)_k} \int_{-1}^{1} \left(\frac{d^k}{dx^k}(1-x^2)^{k+\lambda-\frac{1}{2}}\right) p(x)dx \\ &= \frac{(k+\lambda)\Gamma(\lambda)}{(-2)^k\sqrt{\pi}\Gamma(k+\lambda+\frac{1}{2})} \int_{-1}^{1} \left(\frac{d^k}{dx^k}(1-x^2)^{k+\lambda-\frac{1}{2}}\right) p(x)dx. \end{aligned} \tag{2.3}$$

Therefore, by (2.3), we obtain the following proposition.

Proposition 2.1. *For $p(x) \in \mathbb{P}_n$, let*

$$p(x) = \sum_{n=0}^{n} d_k C_k^{(\lambda)}(x),\ (d_k \in \mathbb{R}).$$

Then

$$d_k = \frac{(k+\lambda)\Gamma(\lambda)}{(-2)^k\sqrt{\pi}\Gamma(k+\lambda+\frac{1}{2})} \int_{-1}^{1} \left(\frac{d^k}{dx^k}(1-x^2)^{k+\lambda-\frac{1}{2}}\right) p(x)dx.$$

For example, let $p(x) = x^n \in \mathbb{P}_n$. From Proposition 2.1, we note that

$$\begin{aligned} d_k &= \frac{(k+\lambda)\Gamma(\lambda)}{(-2)^k\sqrt{\pi}\Gamma(k+\lambda+\frac{1}{2})} \int_{-1}^{1} \left(\frac{d^k}{dx^k}(1-x^2)^{k+\lambda-\frac{1}{2}}\right) x^n dx \\ &= (-n) \int_{-1}^{1} \left(\frac{d^{k-1}}{dx^{k-1}}(1-x^2)^{k+\lambda-\frac{1}{2}}\right) x^{n-1} dx \times \left(\frac{(k+\lambda)\Gamma(\lambda)}{\sqrt{\pi}(-2)^k\Gamma\left(k+\lambda+\frac{1}{2}\right)}\right) \\ &= \cdots \\ &= \frac{(k+\lambda)n!\Gamma(\lambda)}{(n-k)!2^k\sqrt{\pi}(k+\frac{1}{2}+\lambda)} \int_{-1}^{1} (1-x^2)^{k+\lambda-\frac{1}{2}} x^{n-k} dx \\ &= \left(1+(-1)^{n-k}\right) \frac{(k+\lambda)n!\Gamma(\lambda)}{(n-k)!2^k\sqrt{\pi}\Gamma(k+\frac{1}{2}+\lambda)} \int_{0}^{1} (1-x^2)^{k+\lambda-\frac{1}{2}} x^{n-k} dx. \end{aligned} \tag{2.4}$$

Let us assume that $n-k\equiv 0 \pmod 2$. Then, by (2.4), we get

$$\begin{aligned} d_k &= \frac{(k+\lambda)n!\Gamma(\lambda)}{(n-k)!2^k\sqrt{\pi}\Gamma(k+\frac{1}{2}+\lambda)} B\left(k+\lambda+\frac{1}{2}, \frac{n-k+1}{2}\right) \\ &= \frac{\Gamma\left(\frac{n-k+1}{2}\right)\Gamma\left(k+\lambda+\frac{1}{2}\right)}{\Gamma\left(\frac{n+k+2\lambda+2}{2}\right)}, \end{aligned} \tag{2.5}$$

where $B(\alpha,\beta)$ is the beta function which is defined by $B(\alpha,\beta)=\frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$.

It is easy to show that

$$\begin{aligned} \Gamma\left(\frac{n-k+1}{2}\right) &= \frac{n-k-1}{2}\Gamma\left(\frac{n-k-1}{2}\right) \\ &= \left(\frac{n-k-1}{2}\right)\left(\frac{n-k-3}{2}\right)\Gamma\left(\frac{n-k-3}{2}\right) = \cdots \\ &= \frac{\left(\frac{n-k}{2}\right)\left(\frac{n-k-1}{2}\right)\left(\frac{n-k-2}{2}\right)\cdots\frac{2}{2}\Gamma\left(\frac{1}{2}\right)}{\left(\frac{n-k}{2}\right)\left(\frac{n-k-2}{2}\right)\cdots\left(\frac{2}{2}\right)} = \frac{(n-k)!\sqrt{\pi}}{2^{n-k}\left(\frac{n-k}{2}\right)!}. \end{aligned} \tag{2.6}$$

Therefore, by (2.5) and (2.6), we obtain the following identity:

$$x^n = \sum_{0\le k\le n, n-k\equiv 0 \pmod 2} \frac{(k+\lambda)n!\Gamma(\lambda)}{2^n\left(\frac{n-k}{2}\right)!\Gamma\left(\frac{n+k+2\lambda+2}{2}\right)} C_k^{(\lambda)}(x). \tag{2.7}$$

Let us take $p(x)=B_n(x)\in\mathbb{P}_n$. Then, by (1.10), we get

$$\begin{aligned} d_k &= \frac{(k+\lambda)\Gamma(\lambda)}{(-2)^k\sqrt{\pi}\Gamma\left(k+\lambda+\frac{1}{2}\right)}\int_{-1}^{1}\left(\frac{d^k}{dx^k}(1-x^2)^{k+\lambda-\frac{1}{2}}\right)B_n(x)dx \\ &= \frac{(k+\lambda)\Gamma(\lambda)(-n)}{(-2)^k\sqrt{\pi}\Gamma\left(k+\lambda+\frac{1}{2}\right)}\int_{-1}^{1}\left(\frac{d^{k-1}}{dx^{k-1}}(1-x^2)^{k+\lambda-\frac{1}{2}}\right)B_{n-1}(x)dx = \cdots \\ &= \frac{(k+\lambda)\Gamma(\lambda)(-n)(-(n-1))\cdots(-(n-k+1))}{(-2)^k\sqrt{\pi}\Gamma\left(k+\lambda+\frac{1}{2}\right)}\int_{-1}^{1}(1-x^2)^{k+\lambda-\frac{1}{2}}B_{n-k}(x)dx \\ &= \frac{(k+\lambda)\Gamma(\lambda)}{2^k\sqrt{\pi}\Gamma(k+\lambda+\frac{1}{2})}\times\frac{n!}{(n-k)!}\int_{-1}^{1}(1-x^2)^{k+\lambda-\frac{1}{2}}B_{n-k}(x)dx. \end{aligned} \tag{2.8}$$

From (1.10) and (2.7), we can derive the following equation:

$$\begin{aligned} \int_{-1}^{1}(1-x^2)^{k+\lambda-\frac{1}{2}}B_{n-k}(x)dx &= \sum_{l=0}^{n-k}\binom{n-k}{l}B_{n-k-l}\int_{-1}^{1}(1-x^2)^{k+\lambda-\frac{1}{2}}x^l dx \\ &= \sum_{l=0}^{n-k}\binom{n-k}{l}B_{n-k-l}(1+(-1)^l)\int_{0}^{1}(1-x^2)^{k+\lambda-\frac{1}{2}}x^l dx. \end{aligned} \tag{2.9}$$

Let us consider that $l \equiv 0 \pmod 2$. Then, by (2.9), we get

$$\begin{aligned} &\int_{-1}^{1}(1-x^2)^{k+\lambda-\frac{1}{2}}B_{n-k}(x)dx \\ =&2\sum_{0\le l\le n-k,\ l\equiv 0 \pmod 2}\binom{n-k}{l}B_{n-k-l}\int_0^1(1-x^2)^{k+\lambda-\frac{1}{2}}x^l dx \\ =&\sum_{0\le l\le n-k,\ l\equiv 0 \pmod 2}\binom{n-k}{l}B_{n-k-l}\int_0^1(1-y)^{k+\lambda-\frac{1}{2}}y^{\frac{l-1}{2}}dy \\ =&\sum_{0\le l\le n-k,\ l\equiv 0 \pmod 2}\binom{n-k}{l}B_{n-k-l}\frac{\Gamma\left(k+\lambda+\frac{1}{2}\right)\Gamma\left(\frac{l+1}{2}\right)}{\Gamma\left(\frac{2k+2\lambda+l+2}{2}\right)}. \end{aligned} \tag{2.10}$$

For $l \in \mathbb{Z}_+$ with $l \equiv 0$ mod 2), we have

$$\begin{aligned} \Gamma\left(\frac{l+1}{2}\right) &= \Gamma\left(\frac{l-1}{2}+1\right) = \frac{l-1}{2}\Gamma\left(\frac{l-1}{2}\right) \\ &= \left(\frac{l-1}{2}\right)\left(\frac{l-3}{2}\right)\Gamma\left(\frac{l-3}{2}\right) = \cdots \\ &= \left(\frac{l-1}{2}\right)\left(\frac{l-3}{2}\right)\cdots\left(\frac{1}{2}\right)\Gamma\left(\frac{1}{2}\right) = \frac{\left(\frac{1}{2}\right)^l l!\Gamma\left(\frac{1}{2}\right)}{\left(\frac{l}{2}\right)!} = \frac{l!\sqrt{\pi}}{2^l\left(\frac{l}{2}\right)!}. \end{aligned} \tag{2.11}$$

By (2.10) and (2.11), we get

$$\begin{aligned} &\int_{-1}^{1}(1-x^2)^{k+\lambda-\frac{1}{2}}B_{n-k}(x)dx \\ =&\sum_{0\le l\le n-k,\ l\equiv 0 \pmod 2}\binom{n-k}{l}B_{n-k-l}\frac{\Gamma\left(k+\lambda+\frac{1}{2}\right)\Gamma\left(\frac{l+1}{2}\right)}{\Gamma\left(\frac{2k+2\lambda+l+2}{2}\right)} \\ =&\sum_{0\le l\le n-k,\ l\equiv 0 \pmod 2}\binom{n-k}{l}B_{n-k-l}\frac{l!\sqrt{\pi}}{2^l\left(\frac{l}{2}\right)!}\times\frac{\Gamma\left(k+\lambda+\frac{1}{2}\right)}{\Gamma\left(\frac{2k+2\lambda+l+2}{2}\right)}. \end{aligned} \tag{2.12}$$

From (2.8) and (2.12), we have

$$d_k = \frac{n!(k+\lambda)\Gamma(\lambda)}{2^k(n-k)!}\sum_{0\le l\le n-k,\ l\equiv 0 \pmod 2}\frac{\binom{n-k}{l}B_{n-k-l}l!}{2^l\left(\frac{l}{2}\right)!\Gamma\left(\frac{2k+2\lambda+l+2}{2}\right)}. \tag{2.13}$$

Therefore, by (2.13) and Proposition 2.1, we obtain the following theorem.

Theorem 2.2. *For $n \in \mathbb{Z}_+$, we have*

$$\frac{B_n(x)}{n!} = \Gamma(\lambda)\sum_{k=0}^{n}\left(\frac{(k+\lambda)}{2^k(n-k)!}\sum_{0\le l\le n-k,\ l\equiv 0 \ (mod\ 2)}\frac{\binom{n-k}{l}B_{n-k-l}l!}{2^l\left(\frac{l}{2}\right)!\Gamma\left(\frac{2k+2\lambda+l+2}{2}\right)}\right)C_k^{(\lambda)}(x).$$

By the same method, we get

$$\frac{E_n(x)}{n!} = \Gamma(\lambda)\sum_{k=0}^{n}\left(\frac{(k+\lambda)}{2^k(n-k)!}\sum_{0\le l\le n-k,\ l\equiv 0 \pmod 2}\frac{\binom{n-k}{l}E_{n-k-l}l!}{2^l\left(\frac{l}{2}\right)!\Gamma\left(\frac{2k+2\lambda+l+2}{2}\right)}\right)C_k^{(\lambda)}(x). \tag{2.14}$$

From (1.1), we note that

$$\begin{aligned}
&C_{n-k}^{(\lambda)}(x)C_k^{(\lambda)}(x)\\
=&\binom{n-k+2\lambda-1}{n-k}\sum_{l=0}^{n-k}\frac{\binom{n-k}{l}(2\lambda+n-k)_l}{\left(\lambda+\frac{1}{2}\right)_l}\left(\frac{x-1}{2}\right)^l\binom{k+2\lambda-1}{k}\sum_{m-0}^{k}\frac{\binom{k}{m}(2\lambda+k)_m}{\left(\lambda+\frac{1}{2}\right)_m}\left(\frac{x-1}{2}\right)^m\\
=&\binom{n-k+2\lambda-1}{n-k}\binom{k+2\lambda-1}{k}\sum_{p=0}^{n}\left(\sum_{m=0}^{p}\frac{\binom{n-k}{p-m}\binom{k}{m}(2\lambda+k)_m(2\lambda+n-k)_{p-m}}{\left(\lambda+\frac{1}{2}\right)_m\left(\lambda+\frac{1}{2}\right)_{p-m}}\right)\left(\frac{x-1}{2}\right)^p.
\end{aligned}\tag{2.15}$$

Let us take $p(x)=C_k^{(\lambda)}(x)C_{n-k}^{(\lambda)}(x)\in\mathbb{P}_n$. From Proposition 2.1, $p(x)$ can be rewritten as

$$p(x)=C_k^{(\lambda)}(x)C_{n-k}^{(\lambda)}(x)=\sum_{r=0}^{n}d_rC_r^{(\lambda)}(x),\ (d_r\in\mathbb{R}).\tag{2.16}$$

Then, by Proposition 2.1 and (2.15), we get

$$\begin{aligned}
d_r=&\frac{(r+\lambda)\Gamma(\lambda)}{(-2)^r\sqrt{\pi}\Gamma\left(r+\lambda+\frac{1}{2}\right)}\int_{-1}^{1}\left(\frac{d^r}{dx^r}(1-x^2)^{r+\lambda-\frac{1}{2}}\right)C_k^{(\lambda)}(x)C_{n-k}^{(\lambda)}(x)dx\\
=&\frac{(r+\lambda)\Gamma(\lambda)}{(-2)^r\sqrt{\pi}\Gamma\left(r+\lambda+\frac{1}{2}\right)}\binom{n-k+2\lambda-1}{n-k}\binom{k+2\lambda-1}{k}\\
&\times\sum_{p=0}^{n}\left(\sum_{m=0}^{p}\frac{\binom{n-k}{p-m}\binom{k}{m}(2\lambda+k)_m}{\left(\lambda+\frac{1}{2}\right)_m\left(\lambda+\frac{1}{2}\right)_{p-m}}(2\lambda+n-k)_{p-m}\right)\\
&\times\int_{-1}^{1}\left(\frac{d^r}{dx^r}(1-x^2)^{r+\lambda-\frac{1}{2}}\right)\left(\frac{x-1}{2}\right)^p dx\\
=&\frac{(r+\lambda)\Gamma(\lambda)}{(-2)^r\sqrt{\pi}\Gamma\left(r+\lambda+\frac{1}{2}\right)}\binom{n-k+2\lambda-1}{n-k}\binom{k+2\lambda-1}{k}\\
&\times\sum_{p=r}^{n}\left(\sum_{m=0}^{p}\frac{\binom{n-k}{p-m}\binom{k}{m}(2\lambda+k)_m}{\left(\lambda+\frac{1}{2}\right)_m\left(\lambda+\frac{1}{2}\right)_{p-m}}(2\lambda+n-k)_{p-m}\right)\\
&\times\int_{-1}^{1}\left(\frac{d^r}{dx^r}(1-x^2)^{r+\lambda-1}\right)\left(\frac{x-1}{2}\right)^p dx
\end{aligned}\tag{2.17}$$

It is not difficult to show that

$$\begin{aligned}
&\int_{-1}^{1}\left(\frac{d^r}{dx^r}(1-x^2)^{r+\lambda-\frac{1}{2}}\right)\left(\frac{x-1}{2}\right)^p dx=\frac{(-1)^rp!}{2^p(p-r)!}\int_{-1}^{1}(1-x^2)^{r+\lambda-\frac{1}{2}}(1-x)^{p-r}(-1)^{p-r}dx\\
=&\frac{(-1)^pp!}{2^p(p-r)!}\int_{-1}^{1}(1-x)^{p+\lambda-\frac{1}{2}}(1+x)^{r+\lambda-\frac{1}{2}}dx=\frac{(-1)^pp!}{2^p(p-r)!}\int_{0}^{1}(2-2y)^{p+\lambda-\frac{1}{2}}(2y)^{r+\lambda-\frac{1}{2}}2dy\\
=&\frac{(-1)^p2^{p+\lambda-\frac{1}{2}+r+\lambda-\frac{1}{2}+1}}{2^p}\times\frac{p!}{(p-r)!}\int_{0}^{1}(1-y)^{p+\lambda-\frac{1}{2}}y^{r+\lambda-\frac{1}{2}}dy\\
=&(-1)^p2^{r+2\lambda}\frac{p!}{(p-r)!}\times\frac{\Gamma\left(p+\lambda+\frac{1}{2}\right)\Gamma\left(r+\lambda+\frac{1}{2}\right)}{\Gamma\left(r+p+2\lambda+1\right)}.
\end{aligned}\tag{2.18}$$

From the fundamental theorem of gamma function, we have

$$\begin{aligned}\frac{\Gamma\left(p+\lambda+\frac{1}{2}\right)}{\Gamma(r+p+2\lambda+1)} &= \frac{\left(p+\lambda-\frac{1}{2}\right)\cdots\left(\lambda+\frac{1}{2}\right)\Gamma\left(\lambda+\frac{1}{2}\right)}{(r+p+2\lambda)\cdots 2\lambda\Gamma(2\lambda)} \\ &= \frac{\left(\lambda+\frac{1}{2}\right)_p\sqrt{\pi}2^{1-\lambda}}{(2\lambda)_{r+p+1}\Gamma(\lambda)}.\end{aligned} \tag{2.19}$$

By (2.18) and (2.19), we get

$$\begin{aligned}&\int_{-1}^{1}\left(\frac{d^r}{dx^r}(1-x^2)^{r+\lambda-\frac{1}{2}}\right)\left(\frac{x-1}{2}\right)^p dx \\ =&(-1)^p 2^{r+2\lambda}\frac{p!}{(p-r)!}\times\frac{\Gamma\left(p+\lambda+\frac{1}{2}\right)\Gamma\left(r+\lambda+\frac{1}{2}\right)}{\Gamma(r+p+2\lambda+1)} \\ =&(-1)^p 2^{r+2\lambda}\frac{p!}{(p-r)!}\Gamma\left(r+\lambda+\frac{1}{2}\right)\times\frac{\left(\lambda+\frac{1}{2}\right)_p\sqrt{\pi}2^{1-\lambda}}{(2\lambda)_{r+p+1}\Gamma(\lambda)} \\ =&(-1)^p 2^{r+\lambda+1}\frac{p!}{(p-r)!}\Gamma\left(r+\lambda+\frac{1}{2}\right)\times\frac{\left(\lambda+\frac{1}{2}\right)_p\sqrt{\pi}}{(2\lambda)_{r+p+1}\Gamma(\lambda)}.\end{aligned} \tag{2.20}$$

From (2.17) and (2.20), we have

$$\begin{aligned}d_r =&\frac{(r+\lambda)\Gamma(\lambda)}{(-2)^r\sqrt{\pi}\Gamma\left(r+\lambda+\frac{1}{2}\right)}\binom{n-k+2\lambda-1}{n-k}\binom{k+2\lambda-1}{k} \\ &\times\sum_{p=r}^{n}\left(\sum_{m=0}^{p}\frac{\binom{n-k}{p-m}\binom{k}{m}(2\lambda+k)_m}{\left(\lambda+\frac{1}{2}\right)_m\left(\lambda+\frac{1}{2}\right)_{p-m}}(2\lambda+n-k)_{p-m}\right) \\ &\times(-1)^p 2^{r+\lambda+1}\frac{p!}{(p-r)!}\Gamma\left(\lambda+\frac{1}{2}+r\right)\times\frac{\left(\lambda+\frac{1}{2}\right)_p\sqrt{\pi}}{(2\lambda)_{r+p+1}\Gamma(\lambda)} \\ =&(-1)^{r+p}2^{\lambda+1}(r+\lambda)\binom{n-k+2\lambda-1}{n-k}\binom{k+2\lambda-1}{k} \\ &\times\sum_{p=r}^{n}\left(\sum_{m=0}^{p}\frac{\binom{n-k}{p-m}\binom{k}{m}(2\lambda+k)_m}{\left(\lambda+\frac{1}{2}\right)_m\left(\lambda+\frac{1}{2}\right)_{p-m}}(2\lambda+n-k)_{p-m}\frac{p!\left(\lambda+\frac{1}{2}\right)_p}{(p-r)!(2\lambda)_{r+p+1}}\right).\end{aligned} \tag{2.21}$$

Therefore, by (2.21), we obtain the following theorem.

Theorem 2.3. *For $n, k \in \mathbb{Z}_+$ with $n \geq k$, we have*

$$\begin{aligned}&C_{n-k}^{(\lambda)}(x)C_k^{(\lambda)}(x) \\ &= 2^{\lambda+1}\binom{n-k+2\lambda-1}{n-k}\binom{k+2\lambda-1}{k}\sum_{r=0}^{n}\sum_{p=r}^{n}\sum_{m=0}^{p}\left\{(r+\lambda)(-1)^{p+r}\times\right. \\ &\left.\frac{\binom{n-k}{p-m}\binom{k}{m}(2\lambda+k)^m(2\lambda+n-k)_{p-m}p!\left(\lambda+\frac{1}{2}\right)_p}{\left(\lambda+\frac{1}{2}\right)_m\left(\lambda+\frac{1}{2}\right)_{p-m}(p-r)!(2\lambda)_{r+p+1}}\right\}C_r^{(\lambda)}(x).\end{aligned}$$

Let us take $p(x)=C_n^{(\lambda)}(x)\in\mathbb{P}_n$. Then, from (1.1), we have

$$\begin{aligned}C_n^{(\lambda)}(x)&=\frac{\Gamma\left(\lambda+\frac{1}{2}\right)\Gamma(n+2\lambda)}{\Gamma(2\lambda)\Gamma\left(n+\lambda+\frac{1}{2}\right)}P_n^{\left(\lambda-\frac{1}{2},\lambda-\frac{1}{2}\right)}(x)\\&=\frac{(n+2\lambda-1)\cdots(2\lambda)}{\left(n+\lambda-\frac{1}{2}\right)\cdots\left(\lambda+\frac{1}{2}\right)}P_n^{\left(\lambda-\frac{1}{2},\lambda-\frac{1}{2}\right)}(x)=\frac{\binom{n+2\lambda-1}{n}}{\binom{n+\lambda-\frac{1}{2}}{n}}P_n^{\left(\lambda-\frac{1}{2},\lambda-\frac{1}{2}\right)}(x).\end{aligned}\tag{2.22}$$

In a previous paper, we have shown that

$$P_n^{(\alpha,\beta)}(x)=\sum_{k=0}^{n}\binom{n+\alpha}{n-k}\binom{n+\beta}{k}\left(\frac{x-1}{2}\right)^k\left(\frac{x+1}{2}\right)^{n-k},\ \text{(see [11])}.\tag{2.23}$$

From (2.22) and (2.23), we have

$$C_n^{(\lambda)}(x)=\frac{\binom{n+2\lambda-1}{n}}{\binom{n+\lambda-\frac{1}{2}}{n}}\sum_{k=0}^{n}\binom{n+\lambda-\frac{1}{2}}{n-k}\binom{n+\lambda-\frac{1}{2}}{k}\left(\frac{x-1}{2}\right)^k\left(\frac{x+1}{2}\right)^{n-k},\tag{2.24}$$

and

$$\frac{d^k}{dx^k}C_n^{(\lambda)}(x)=2^k\lambda^kC_{n-k}^{(\lambda+k)}(x).\tag{2.25}$$

Let $p(x)=C_n^{(\lambda)}(x)=\sum_{k=0}^{n}d_kC_k^{(\lambda)}(x)$. Then, by Proposition 2.1, we get

$$\begin{aligned}d_k&=\frac{(k+\lambda)\Gamma(\lambda)}{(-2)^k\sqrt{\pi}\Gamma\left(k+\lambda+\frac{1}{2}\right)}\int_{-1}^{1}\left(\frac{d^k}{dx^k}(1-x^2)^{k+\lambda-\frac{1}{2}}\right)C_n^{(\lambda)}(x)dx\\&=\frac{(k+\lambda)\Gamma(\lambda)}{(-2)^k\sqrt{\pi}\Gamma\left(k+\lambda+\frac{1}{2}\right)}(-1)^k2^k\lambda^k\int_{-1}^{1}(1-x^2)^{k+\lambda-\frac{1}{2}}C_{n-k}^{(\lambda+k)}(x)dx\\&=\frac{\lambda^k(k+\lambda)\Gamma(\lambda)}{\sqrt{\pi}\Gamma\left(k+\lambda+\frac{1}{2}\right)}\int_{-1}^{1}(1-x^2)^{k+\lambda-\frac{1}{2}}C_{n-k}^{(\lambda+k)}(x)dx.\end{aligned}\tag{2.26}$$

By (2.24), we get

$$\begin{aligned}&C_{n-k}^{(\lambda+k)}(x)\\=&\frac{\binom{n-k+2(\lambda+k)-1}{n-k}}{\binom{n-k+\lambda+k-\frac{1}{2}}{n-k}}\sum_{l=0}^{n-k}\binom{n-k+\lambda+k-\frac{1}{2}}{n-k-l}\binom{n-k+\lambda+k-\frac{1}{2}}{l}\left(\frac{x-1}{2}\right)^l\left(\frac{x+1}{2}\right)^{n-k-l}\\=&\frac{\binom{n+k+2\lambda-1}{n-k}}{\binom{n+\lambda-\frac{1}{2}}{n-k}}\sum_{l=0}^{n-k}\binom{n+\lambda-\frac{1}{2}}{n-k-l}\binom{n+\lambda-\frac{1}{2}}{l}\left(\frac{x-1}{2}\right)^l\left(\frac{x+1}{2}\right)^{n-k-l}.\end{aligned}\tag{2.27}$$

From (2.26) and (2.27), we have

$$\begin{aligned}d_k=&\frac{\lambda^k(k+\lambda)\Gamma(\lambda)}{\sqrt{\pi}\Gamma\left(k+\lambda+\frac{1}{2}\right)}\times\frac{\binom{n+k+2\lambda-1}{n-k}}{\binom{n+\lambda-\frac{1}{2}}{n-k}}\sum_{l=0}^{n-k}\binom{n+\lambda-\frac{1}{2}}{n-k-l}\binom{n+\lambda-\frac{1}{2}}{l}(-1)^l\left(\frac{1}{2}\right)^{n-k}\\&\times\int_{-1}^{1}(1-x)^{k+\lambda-\frac{1}{2}+l}(1+x)^{\lambda+n-\frac{1}{2}-l}dx.\end{aligned}\tag{2.28}$$

It is easy to show that

$$\begin{aligned}&\int_{-1}^{1}(1-x)^{k+\lambda-\frac{1}{2}+l}(1+x)^{\lambda+n-l-\frac{1}{2}}dx=\int_{0}^{1}(2-2y)^{k+\lambda-\frac{1}{2}+l}(2y)^{\lambda+n-l-\frac{1}{2}}2dy\\ =&2^{n+2\lambda+k}\int_{0}^{1}(1-y)^{k+\lambda+l-\frac{1}{2}}y^{\lambda+n-l-\frac{1}{2}}dy\\ =&2^{k+n+2\lambda}\frac{\Gamma\left(k+\lambda+l+\frac{1}{2}\right)\Gamma\left(\lambda+n-l+\frac{1}{2}\right)}{\Gamma(k+n+2\lambda)}.\end{aligned} \tag{2.29}$$

By fundamental theorem of gamma function, we see that

$$\Gamma\left(k+\lambda+l+\frac{1}{2}\right)=\binom{k+\lambda+l-\frac{1}{2}}{l}l!\Gamma\left(k+\lambda+\frac{1}{2}\right), \tag{2.30}$$

$$\Gamma\left(\lambda+n-l+\frac{1}{2}\right)=\binom{\lambda+n-l-\frac{1}{2}}{n-l}(n-l)!\Gamma\left(\lambda+\frac{1}{2}\right), \tag{2.31}$$

and

$$\Gamma\left(k+2\lambda+n\right)=\binom{k+2\lambda+n-1}{n+k}(n+k)!\Gamma(2\lambda). \tag{2.32}$$

As is well known, the duplication formula for gamma function is given by

$$\Gamma(z)\Gamma\left(z+\frac{1}{2}\right)=2^{1-2z}\sqrt{\pi}\Gamma(2z). \tag{2.33}$$

By (2.29), (2.30), (2.31) and (2.32), we get

$$\begin{aligned}&\int_{-1}^{1}(1-x)^{k+\lambda+l-\frac{1}{2}}(1+x)^{\lambda+n-l-\frac{1}{2}}dx\\ =&2^{k+n+1}\frac{\binom{k+\lambda+l-\frac{1}{2}}{l}\binom{\lambda+n-l-\frac{1}{2}}{n-l}\Gamma\left(k+\lambda+\frac{1}{2}\right)}{\binom{n}{l}\binom{k+2\lambda+n-1}{n+k}\binom{n+k}{k}k!\Gamma(\lambda)}\sqrt{\pi}.\end{aligned} \tag{2.34}$$

From (2.28) and (2.34), we have

$$\begin{aligned}d_k=&\lambda^k(k+\lambda)2^{2k+1}\frac{\binom{n+k+2\lambda-1}{n-k}}{\binom{n+\lambda-\frac{1}{2}}{n-k}}\\ &\times\sum_{l=0}^{n-k}\binom{n+\lambda-\frac{1}{2}}{n-k-l}\binom{n+\lambda-\frac{1}{2}}{l}(-1)^l\frac{\binom{k+\lambda+l-\frac{1}{2}}{l}\binom{\lambda+n-l-\frac{1}{2}}{n-l}}{\binom{n}{l}\binom{k+2\lambda+n-1}{n+k}\binom{n+k}{k}k!}.\end{aligned} \tag{2.35}$$

Therefore, by (2.35), we obtain the following theorem.

Theorem 2.4. *For $n \in \mathbb{Z}_+$, we have*

$$\begin{aligned}C_n^{(\lambda)}(x)=&\sum_{k=0}^{n}\left\{\frac{\lambda^k(k+\lambda)2^{2k+1}\binom{n+k+2\lambda-1}{n-k}}{\binom{n+\lambda-\frac{1}{2}}{n-k}}\right.\\ &\left.\times\sum_{l=0}^{n-k}\frac{\binom{n+\lambda-\frac{1}{2}}{n-k-l}\binom{n+\lambda-\frac{1}{2}}{l}(-1)^l\binom{k+\lambda+l-\frac{1}{2}}{l}}{\binom{n}{l}\binom{k+2\lambda+n-1}{n+k}\binom{n+k}{k}k!}\binom{\lambda+n-l-\frac{1}{2}}{n-l}\right\}C_k^{(\lambda)}(x).\end{aligned}$$

References

[1] W. A. Al-Salam, L. Carlitz, *The Gegenbauer addition theorem*, J. Math. and Phys. **42(1963)**, 147-156.

[2] A. Bayad and T. Kim, *Identities involving values of Bernstein, q-Bernoulli, and q-Euler polynomials*, Russ. J. Math. Phys. **18(2011)**, no. 2, 133-143.

[3] A. Bayad, *Modular properties of elliptic Bernoulli and Euler functions*, Adv. Stud. Contemp. Math. **20(2010)**, no. 3, 389-401.

[4] I. N. Cangul, V. Kurt, H. Ozden and Y. Simsek, *On the higher-order w-q-Genocchi numbers*, Adv. Stud. Contemp. Math. **19(2009)**, no. 1, 39-57.

[5] L.-C. Jang, *A family of Barnes-type multiple twisted q-Euler numbers and polynomials related to Fermionic p-adic invariant integrals on* $\mathbb{Z}_p$, J. Comput. Anal. Appl. **13(2011)**, no. 2, 376-387.

[6] S. Khan, A. A, Al-Gonah, G. Yasmin, *Generalized and mixed type Gegenbauer polynomials*, J. Math. Anal. Appl. **390 (2012)**, no. 1, 197-207.

[7] T. Kim, J. Choi, Y. H. Kim and C. S. Ryoo, *On q-Bernstein and q-Hermite polynomials*, Proc. Jangjeon Math. Soc. **14 (2011)**, no. 2, 215-221.

[8] T. Kim, *Some identities on the q-Euler polynomials of higher order and q-Stirling numbers by the fermionic p-adic integral on* $\mathbb{Z}_p$, Russ. J. Math. Phys. **16(2009)**, no. 4, 484-491.

[9] T. Kim, *Symmetry of power sum polynomials and multivariate fermionic p-adic invariant integral on* $\mathbb{Z}_p$, Russ. J. Math. Phys. **16(2009)**, no. 1, 93-96.

[10] T. Kim, *Power series and asymptotic series associated with the q-analog of the two-variable p-adic L-function*, Russ. J. Math. Phys. **12(2005)**, no. 2, 186-196.

[11] D. S. Kim, T. Kim and D. V. Dolgy, *Some Identities on Bernoulli and Hermite Polynomials Associated with Jacobi Polynomials* (submitted).

[12] H. Y. Lee, N. S. Jung, J. Y. Kang and C. S. Ryoo, *Some identities on the higher-order-twisted q-Euler numbers and polynomials with weight* α, Adv. Difference Equ. **2012**, 2012:21.

[13] J. A. McFadden, *A diagonal expansion in Gegenbauer polynomials for a class of second-order probability densities*, SIAM J. Appl. Math. **14(1966)**, 1433-1436.

[14] H. Ozden, Cangul and Y. Simsek, *Remarks on q-Bernoulli numbers associated with Daehee numbers*, Adv. Stud. Contemp. Math. **18(2009)**, no. 1, 41-48.

[15] S.-H. Rim, S-J. Lee, *Some identities on the twisted* (h,q)*-Genocchi numbers and polynomials associated with q-Bernstein polynomials*, Int. J. Math. Math. Sci. 2011, Art. ID 482840, 8 pp.

[16] C. S. Ryoo, *Some relations between twisted q-Euler numbers and Bernstein polynomials*, Adv. Stud. Contemp. Math. **21(2011)**, no. 2, 217-223.

[17] C.S. Ryoo, *On the generalized Barnes type multiple q-Euler polynomials twisted by ramified roots of unity*, Proc. Jangjeon Math. Soc. **13(2010)**, no. 2, 255-263.

[18] M. Shah, *Applications of Gegenbauer (ultraspherical) polynomials in cooling of a heated cylinder*, An. Univ. Timisoara Ser. Sti. Mat. **8(1970)**, 207-212.

[19] M. Shah, *Some properties associated with generalized integral transform in two variables*, Univ. Brasov Lucrari Stiint, **18(1976)**, 15-27.

[20] Y. Simsek, *Special functions related to Dedekind-type DC-sums and their applications*, Russ. J. Math. Phys. **17(2010)**, no. 4, 495-508.

[21] Y. Simsek, *Interpolation function of generalized q-Bernstein-type basis polynomials and applications, Curves and surfaces*, 647-662, Lecture Notes in Comput. Sci., 6920,Springer, Heidelberg, 2012.

[22] Y. Simsek, *Generating functions of the twisted Bernoulli numbers and polynomials associated with their interpolation functions*, Adv. Stud. Contemp. Math. **16(2008)**, no. 2, 251-278.

[23] I. N. Sneddon, *The evaluation of an integral involving the product of two Gegenbauer polynomials*, SIAM Rev. **9(1967)** 569-572.

[1] Department of Mathematics, Sogang University, Seoul 121-742, Republic of Korea
E-mail address: `dskim@sogang.ac.kr`

[2] Department of Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea.
E-mail address: `tkkim@kw.ac.kr`

[3] Department of Mathematics Education, Kyungpook National University, Taegu 702-701, Republic of Korea.
E-mail address: `shrim@knu.ac.kr`